



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Preface: The mountain cryosphere - a holistic view on processes and their interactions

Gruber, Stephan ; Egli, Markus ; Gärtner-Roer, Isabelle ; Hoelzle, Martin

DOI: <https://doi.org/10.1111/j.1468-0459.2012.00468.x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-66969>

Journal Article

Accepted Version

Originally published at:

Gruber, Stephan; Egli, Markus; Gärtner-Roer, Isabelle; Hoelzle, Martin (2012). Preface: The mountain cryosphere - a holistic view on processes and their interactions. *Geografiska Annaler. Series A. Physical Geography*, 94(2):177-182.

DOI: <https://doi.org/10.1111/j.1468-0459.2012.00468.x>

PREFACE:

THE MOUNTAIN CRYOSPHERE – A HOLISTIC VIEW ON PROCESSES AND THEIR INTERACTIONS

STEPHAN GRUBER¹, MARKUS EGLI¹, ISABELLE GÄRTNER-ROER¹ and MARTIN HOELZLE²

¹Department of Geography, University of Zürich, Zürich, Switzerland

²Department of Geosciences, University of Fribourg, Fribourg, Switzerland

Gruber, S., Egli, M., Gärtner-Roer, I. and Hoelzle, M., 2012. Preface: The mountain cryosphere – a holistic view on processes and their interactions. *Geografiska Annaler: Series A, Physical Geography*, **93**, 1–3. doi:10.1111/j.1468-0459.2012.00468.x

This thematic issue of *Geografiska Annaler: Series A, Physical Geography* is dedicated to Professor Wilfried Haeblerli on the occasion of his retirement in January 2012. Paraphrasing the wide and integrative scope of his scientific work, it is titled ‘The mountain cryosphere – A holistic view on processes and their interactions’ and contains contributions arising from keynote papers of a symposium held in Zurich with the same name on 20 January 2012.

The scientific work of Wilfried Haeblerli (Fig. 1) began with his PhD at the University of Basel, Switzerland. After this, Wilfried Haeblerli started to work at the Laboratory of Hydraulics, Hydrology and Glaciology (VAW) at the Swiss Federal Institute of Technology Zurich (ETHZ) in 1974 and became head of its Glaciology Section in 1988. In 1978/1979 he worked with Mark Meier as visiting scientist at the US Geological Survey in Tacoma, Washington and Fairbanks, Alaska. From 1995 until 2012 Wilfried Haeblerli was Full Professor for Physical Geography at the University of Zurich. The influential figures in his early career include Hans Oeschger, Bruno Messerli, Herfried Hoinkes, and his PhD advisor Dietrich Barsch.

Scientific contribution

In his PhD, Wilfried Haeblerli investigated the distribution of permafrost in an Alpine test area (Flüelapass, Eastern Switzerland). Then a marginal and exotic research topic with only few previous




Fig. 1. Wilfried Haeblerli during an excursion to the northern Patagonian ice field. Photo: 11 September 2009, Amanda Maxwell.

studies, it is much due to his work that mountain permafrost research has now developed into an important field in science and practice. The publication of the BTS (Bottom Temperature of the winter Snow cover) method (Haeblerli 1973) and the so-called rules-of-thumb in his PhD (Haeblerli 1975) provided a basis for the spatial delineation of permafrost occurrence in mountains. Aided by GIS modelling, this also enabled spatial estimates (Keller 1992; Hoelzle *et al.* 1993; Hoelzle and

Haeberli 1995; Haeberli *et al.* 1996) visualized as maps to communicate the topic to a broader audience. The investigation of rock glaciers took him from an initial reconstruction of previous research (Fisch *et al.* 1978) over detailed studies of internal structure (Haeberli 1985a), photogrammetric analyses of surface displacement (Haeberli and Schmid 1988), analyses of material composition (Haeberli *et al.* 1999a) to thermal characteristics (VonderMühll and Haeberli 1990) to a broad and detailed state of the art summary given three decades later (Haeberli *et al.* 2006). Drilling in rock glaciers has been pioneered and promoted (Barsch *et al.* 1979; Haeberli *et al.* 1988; 1998). The interaction with glaciers (Haeberli 1979; 1983; 2005), and more generally the role of permafrost in the conditioning of landforms (Haeberli 1996; Gruber and Haeberli 2007) have been recurring themes. A summary of these four decades of mountain permafrost research is given by Haeberli *et al.* (2010).

In the aftermath of the Mattmark ice avalanche that killed 88 construction workers in 1965, and with the expansion of hydropower schemes and other infrastructure, the assessment and understanding of hazards related to glaciers and permafrost gained importance in Switzerland. Early in his career, Wilfried Haeberli performed corresponding research (Haeberli and Röthlisberger 1976; Haeberli 1977; Haeberli *et al.* 1979, 1989a) that was often tied to consulting mandates. Flooding and debris flows in many areas of Switzerland during the summer of 1987 caused considerable damage (Naef *et al.* 1989; Haeberli *et al.* 1990) and triggered continued interest in the role of permafrost degradation in debris flow conditioning (Zimmermann and Haeberli 1992) that was continued in a national research programme (Haeberli *et al.* 1999b). Together with the beginning interest in permafrost in steep bedrock (Haeberli *et al.* 1997; Wegmann *et al.* 1998) this was one foundation for the EU FP6 research Programme PACE (Harris *et al.* 2009). The expertise on combined assessment of glacier and permafrost hazards has also found important foci outside Switzerland (Haeberli *et al.* 2002, 2004a; Huggel *et al.* 2007).

Early in his career, Wilfried Haeberli also started research on ice temperatures of high-elevation Alpine summits, demonstrating that cold ice, and thus the potential for retrieving ice cores as environmental archives, existed (Haeberli 1976). From this, a continued interest in Alpine paleoglaciology

followed in the directions of ice-core analyses (Oeschger *et al.* 1978; Haeberli *et al.* 1983, 1988) and reconstructions of past glacial environments (Haeberli and Penz 1985; Lister *et al.* 1998). Besides ice cores from cold firm areas, cold miniature ice caps formed by superimposed ice provide a further valuable environmental archive (Haeberli *et al.* 2004b). More recent investigations of firm temperatures (Suter *et al.* 2001) supported the conclusion, that not only many miniature ice caps, but also the high-elevation cold firm areas may be lost as environmental archives during coming decades. The connection between firm and ice temperatures, permafrost and natural hazards is explored in several examples (Haeberli *et al.* 1999b; Hasler *et al.* 2011).

In 1983, Wilfried Haeberli became director of the Permanent Service on the Fluctuations of Glaciers, marking the beginning of his three-decade long contribution to international cryosphere and climate monitoring. As a scientist and in his role as the first director of the World Glacier Monitoring Service established in 1986, he made key contributions to the design and implementation of national (Haeberli *et al.* 1993) and global (Haeberli 1985b; Haeberli *et al.* 1989b, 2007; Brown *et al.* 2001; Paul *et al.* 2009) monitoring strategies and systems, as well as their international coordination (Townshend *et al.* 1995; Haeberli *et al.* 2000). In this, he also continued to promote the interpretation of long-term length change measurements (Hoelzle *et al.* 2003) and the systematic use of satellite observations as spatial complements of point-based mass-balance observations (Kieffer *et al.* 2000; Paul *et al.* 2002; Haeberli 2004). This, and the estimation of high-resolution digital elevation models of glacier beds including the modelling of potential future lakes in deglaciating high-mountain regions (Linsbauer *et al.* 2009; Frey *et al.* 2010) is in many ways based on early analyses of inventory data (Haeberli and Hoelzle 1995) that established methods for the evaluation of large samples of glaciers. As co-author and reviewer, he has been actively involved in the compilation of the first four IPCC assessment reports and was committed to the communication of cryosphere-climate interactions (Haeberli 1994; Fischlin and Haeberli 2008; Haeberli and Zemp 2010; Ajai *et al.* 2011).

During his career, Wilfried Haeberli has also taught and inspired many students and young researchers, and though collaboration and joint research contributed to shaping his scientific field.

Content of this issue

The contributions in this issue concern differing topics related to mountain cryosphere processes and phenomena and outline important new directions for research. In the outlines given below and in the contributions themselves, the relation to the work of Wilfried Haeberli is directly evident.

Mountain glaciers are iconic indicators of climate change but the combined and quantitative use of their length changes as a climate proxy remains challenging. This is because differences in size, topography and environmental conditions lead to variations in the magnitude and timing of the response of individual glaciers to regional forcing. Oerlemans (2012) uses a model for representing glacier length fluctuations with only two parameters that relate to sensitivity and response time. This simplicity enables extracting information on climate forcing and glacier parameters from ensembles of glacier records. Results are encouraging in terms of an agreement between forcings reconstructed from neighbouring glaciers, but some differences to forcing inferred from station data exists.

While permafrost research is an important topic in North America, mountain ranges there have received only marginal interest. Lewkowicz *et al.* (2012) describe a unique set of ground and air temperature measurements and a derived spatial model of permafrost in the southern Yukon and northern British Columbia, Canada. They show how winter inversions, which usually influence air temperature gradients below the tree line in valleys, have a dominating influence on the spatial distribution of permafrost and further complicate modelling in mountains. Based on the transfer functions between ground and air temperatures derived from their measurement network, they also argue that local surface and subsurface conditions add to the strong heterogeneity of ground temperatures and will result in differing patterns of temperature evolution in the future.

Rock glaciers are prominent landforms indicative of mountain permafrost and recently have received increased interest in the context of hazard assessment. Using diverse examples, Springman *et al.* (2012) provide an overview of current process studies aimed at understanding the observed recent acceleration and beginning collapse of many rock glaciers. They describe in-situ and laboratory geotechnical testing and measurements of deformation and temperature in the field as well as combined modelling of mechanical sta-

bility, thermal regime, and water flow. From the studies presented it is evident that broad multidisciplinary teams are needed to advance our understanding of the processes involved in rock-glacier degradation.

In glacierized high-alpine areas, non-temperate glaciers may be unique in holding climate signals linked to past atmospheric changes. Because the Alps are surrounded by the major European anthropogenic emission sources, their glaciers are not only an archive for climatic characteristics but also particularly for man-made changes in atmospheric composition. Wagenbach and Preunkert (2012) discuss results from two major coring areas (located in the summit ranges of Monte Rosa and Mt Blanc massif) which largely differ in their snow accumulation rate, and hence, in their accessible time scales. When investigating Alpine ice cores, the intricate role of seasonality of specific compounds (particularly $\delta^{18}\text{O}$) and snow erosion have to be taken into account. Furthermore, an appropriate dating of e.g., the basal layer, for tracing ice formation and landscape evolution is still a major challenge. The adequate application of dating tools to reconstruct the earliest stages of a glacier is particularly intriguing in view of cold, low-elevation miniature ice-caps.

Besides the cold ice found at highest elevations, the Mont Blanc massif provides many more examples of changing cryosphere systems. Deline *et al.* (2012) provide an overview of current knowledge concerning recent and Holocene changes of glaciers and permafrost, as well as their impact on geomorphic phenomena, and ultimately, natural hazards. In this area that has been well documented photographically due to tourism during the past century, clear indication for increasing rock fall activity related to warm periods on decadal and annual scales exist. A large number of ice-bearing fractures has been observed in the detachment areas of rock falls, further supporting the hypothesis that permafrost degradation is an important element in explaining the cause of this increase. The accelerated glacier retreat in the area led to a strong increase in the debris cover on glaciers, providing a self-retardation effect that has reduced summer melt. The discussed glacial hazards are related to ice fall and effects caused by the outburst of glacial lakes and water pockets. Despite the widespread glacier melt, these hazards are unlikely to significantly decrease during the coming decades.

All the contributions are closely related to Wilfried Haeberli's scientific interest and activities and

are a tribute to the leadership and inspiration given by him during the past four decades – one part of his scientific legacy.

Stephan Gruber, Markus Egli, Isabelle Gärtner-Roer, Department of Geography, University of Zürich, Winterthurerstrasse, 190, 8057 Zürich, Switzerland.
Email: stephan.gruber@geo.uzh.ch

Martin Hoelzle, Department of Geosciences, University of Fribourg, Chemin du Musée 4, 1700 Fribourg, Switzerland.

References

- Ajai, Bengtsson, L., Breashears, D., Crutzen, P.J., Fuzzi, S., Haeberli, W., Immerzeel, W.W., Kaser, G., Kennel, C., Kulkarni, A., Pachauri, R., Painter, T.H., Rabassa, J., Ramanathan, V., Robock, A., Rubbia, C., Russell, L., Sánchez Sorondo, M., Schellnhuber, H.K., Sorooshian, S., Stocker, T.F., Thompson, L.G., Toon, O.B. and Zaelke, D., 2011. *Fate of Mountain Glaciers in the Anthropocene*. A Report by the Working Group Commissioned by the Pontifical Academy of Sciences.
- Barsch, D., Fierz, H. and Haeberli, W., 1979. Shallow core drilling and borehole measurements in the permafrost of an active rock glacier near the Grubengletscher, Wallis, Swiss Alps. *Arctic and Alpine Research*, 11 (2), 215–228.
- Brown, J., Haeberli, W., Barry, R. and Nelson, F.E., 2001. The proposed international permafrost monitoring network and service. In: Paepe, R. and Melnikov, V. (eds), *Permafrost Response on Economic Development, Environmental Security and Natural Resources*. Kluwer Academic Publishers, Netherlands, 601–606.
- Deline, P., Gardent, M., Magnin, F. and Ravel, L., 2012. The morphodynamics of the Mont Blanc massif in a changing cryosphere: a comprehensive review. *Geografiska Annaler: Series A, Physical Geography*. doi:10.1111/j.1468-0459.2012.00467.x
- Fisch, W. sen., Fisch, W. jun. and Haeberli, W., 1978. Electrical D.C. resistivity soundings with long profiles on rock glaciers and moraines in the Alps of Switzerland. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 13 (1/2), 239–260.
- Fischlin, A. and Haeberli, W., 2008. Auch in der Schweiz wirkt sich der Klimawandel zunehmend aus. In: *Das Klima ändert – was nun? Der neue UN-Klimabericht (IPCC 2007) und die wichtigsten Ergebnisse aus der Sicht der Schweiz*. OeCC/scnat, Bern, 21–32.
- Frey, H., Haeberli, W., Linsbauer, A., Huggel, C. and Paul, F., 2010. A multi level strategy for anticipating future glacier lake formation and associated hazard potentials. *Natural Hazards and Earth System Science*, 10, 339–352.
- Gruber, S. and Haeberli, W., 2007. Permafrost in steep bedrock slopes and its temperature-related destabilization following climate change. *Journal of Geophysical Research*, 112, F02S18. doi: 10.1029/2006JF000547
- Haeberli, W., 1973. Die Basis-Temperatur der winterlichen Schneedecke als möglicher Indikator für die Verbreitung von Permafrost in den Alpen. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 9 (1–2), 221–227.
- Haeberli, W., 1975. *Untersuchung zur Verbreitung von Permafrost zwischen Flüelapass und Piz Grialettsch (Graubünden)*. Dissertation der Universität Basel. Mitteilung VAW/ETHZ, 17.
- Haeberli, W., 1976. Eistemperaturen in den Alpen. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 11 (2), 203–220.
- Haeberli, W., 1977. Experience with glacier calving and air bubbling in high Alpine water reservoirs. *Journal of Glaciology*, 19 (81), 589–594.
- Haeberli, W., 1979. Holocene push-moraines in Alpine permafrost. *Geografiska Annaler, Series A: Physical Geography*, 61 (1–2), 43–48.
- Haeberli, W., 1983. Permafrost–glacier relationships in the Swiss Alps today and in the past. *Proceedings of the Fourth International Conference on Permafrost*, 415–420.
- Haeberli, W., 1985a. *Creep of mountain permafrost: internal structure and flow of Alpine rock glaciers*. Mitteilung VAW/ETHZ, 74.
- Haeberli, W., 1985b. Global land ice monitoring: present status and future perspectives. *US Department of Energy Report, DOE/EV60235-1*, 216–231.
- Haeberli, W., 1994. Accelerated glacier and permafrost changes in the Alps. In: Beniston, M., (ed.), *Mountain Environments in Changing Climates*. Routledge, London, 91–107.
- Haeberli, W., 1996. On the morphodynamics of ice/debris-transport systems in cold mountain areas. *Norsk Geografisk Tidsskrift*, 50, 3–9.
- Haeberli, W., 2004. Glaciers and ice caps: historical background and strategies of world-wide monitoring. In: Bamber, J.L. and Payne, A.J., (eds), *Mass Balance of the Cryosphere*. Cambridge University Press, Cambridge, 559–578.
- Haeberli, W., 2005. Climate change and glacial/periglacial geomorphodynamics in the Alps: a challenge of historical dimensions. *Geografia Fisica e Dinamica Quaternaria, Suppl.-Vol. II*, 9–14.
- Haeberli, W. and Hoelzle, M., 1995. Application of inventory data for estimating characteristics of and regional climate-change effects on mountain glaciers: a pilot study with the European Alps. *Annals of Glaciology*, 21, 206–212.
- Haeberli, W. and Penz, U., 1985. An attempt to reconstruct glaciological and climatological characteristics of 18ka BP Ice Age glaciers in and around the Swiss Alps. *Norsk Geografisk Tidsskrift/Zeitschrift für Gletscherkunde und Glazialgeologie*, 21, 351–361.
- Haeberli, W. and Röthlisberger, H., 1976. Beobachtungen zum Mechanismus und zu den Auswirkungen von Kalbungen am Grubengletscher (Saastal, Schweiz). *Zeitschrift für Gletscherkunde und Glazialgeologie*, 11 (2), 221–228.
- Haeberli, W. and Schmid, W., 1988. Aerophotogrammetrical monitoring of rock glaciers. *Proceedings of the Fifth International Conference on Permafrost*, 1, 764–769.
- Haeberli, W. and Zemp, M., 2010. The uncertain future of European glaciers. *Europe's Ecological Backbone: Recognising the True Value of our Mountains. EEA Report 6/2010, Box 6.2*, 90–93.
- Haeberli, W., Alean, J.C., Müller, P. and Funk, M., 1989a. Assessing risks from glacier hazards in high mountain

- regions: some experiences in the Swiss Alps. *Annals of Glaciology*, 13, 96–102.
- Haeblerli, W., Barry, R. and Cihlar, J., 2000. Glacier monitoring within the Global Climate Observing System. *Annals of Glaciology*, 31, 241–246.
- Haeblerli, W., Bösch, H., Scherler, K., Østrem, G. and Wallén, C.C., 1989b. *World Glacier Inventory – Status 1988. A contribution to the Global Environment Monitoring System (GEMS) and the International Hydrological Programme*. IAHS/EMS/UNEP/ UNESCO.
- Haeblerli, W., Frauenfelder, R., Kääb, A. and Wagner S., 2004b. Characteristics and potential climatic significance of “miniature ice caps” (crest- and cornice-type low-altitude ice archives). *Journal of Glaciology*, 50 (168), 129–136.
- Haeblerli, W., Hallet, B., Arenson, L., Elconin, R., Humlum, O., Kääb, A., Kaufmann, V., Ladanyi, B., Matsuoka, N., Springman, S. and VonderMühl, D., 2006. Permafrost creep and rock glacier dynamics. *Permafrost and Periglacial Processes*, 17 (3), 189–214. doi: 10.1002/ppp
- Haeblerli, W., Hoelzle, M., Dousse, J.P., Ehrler, C., Gardaz, J.M., Imhof, M., Keller, F., Kunz, P., Lugon, R. and Reynard, M., 1996. *Simulation der Permafrostverbreitung in den Alpen mit geographischen Informationssystemen*. Arbeitsheft NFP 31, vdfHochschulverlag ETH Zürich.
- Haeblerli, W., Hoelzle, M., Kääb, A., Keller, F., VonderMühl, D. and Wagner, S., 1998. Ten years after drilling through the permafrost of the active rock glacier Murtèl, eastern Swiss Alps: answered questions and new perspectives. *Proceedings of the Seventh International Conference on Permafrost, Yellowknife, Canada, Collection Nordica*, 57, 403–410.
- Haeblerli, W., Hoelzle, M., Keller, F., Schmid, W., VonderMühl, D. and Wagner, S., 1993. Monitoring the long-term evolution of mountain permafrost in the Swiss Alps. *Proceedings of the Sixth International Conference on Permafrost*, 1, 214–219.
- Haeblerli, W., Hoelzle, M., Paul, F. and Zemp, M., 2007. Integrated monitoring of mountain glacier as key indicators of global climate change: the European Alps. *Annals of Glaciology*, 46, 150–160.
- Haeblerli, W., Huder, J., Keusen, H.R., Pika, J. and Röthlisberger, H., 1988. Core drilling through rock-glacier permafrost. *Proceedings of the Fifth International Conference on Permafrost*, 2, 937–942.
- Haeblerli, W., Huggel, C., Kääb, A., Zraggen-Oswald, S., Polkvoj, A., Galushkin, I., Zotikov, I. and Osokin, N., 2004a. The Kolka-Karmadon rock/ice slide of 20 September 2002: an extraordinary event of historical dimensions in North Ossetia, Russian Caucasus. *Journal of Glaciology*, 50 (171), 533–546.
- Haeblerli, W., Iken, A. and Siegenthaler, H., 1979. Glaziologische Aspekte beim Bau der Fernmelde-Mehrweckanlage der PTT auf dem Chli Titlis. Mitteilung VAW/ETHZ, 41, “Festschrift Peter Kasser”, 59–75.
- Haeblerli, W., Kääb, A., Hoelzle, M., Bösch, H., Funk, M., VonderMühl, D. and Keller, F., 1999b. *Eisschwund und Naturkatastrophen im Hochgebirge*. Schlussbericht NFP 31, vdfHochschulverlag ETH Zürich.
- Haeblerli, W., Kääb, A., Paul, F., Chiarle, M., Mortara, G., Mazza, A., Deline, P. and Richardson, S., 2002. A surge-type movement at Ghiacciaio del Belvedere and a developing slope instability in the east face of Monte Rosa, Macugnaga, Italian Alps. *Norsk Geografisk Tidsskrift*, 56, 104–111.
- Haeblerli, W., Kääb, A., Wagner, S., VonderMühl, D., Geissler, P., Haas, J.N., Glatzel-Mattheier, H. and Wagenbach, D., 1999a. Pollen analysis and ¹⁴C age of moss remains in a permafrost core recovered from the active rock glacier Murtèl-Corvatsch, Swiss Alps: geomorphological and glaciological implications. *Journal of Glaciology*, 45 (149), 1–8.
- Haeblerli, W., Noetzli, J., Arenson, L., Delaloye, R., Gärtner-Roer, I., Gruber, S., Isaksen, K., Kneisel, C., Krautblatter, M. and Phillips, M., 2010. Mountain permafrost: Development and challenges of a young research field. *Journal of Glaciology*, 56 (200), 1043–1058.
- Haeblerli, W., Rickenmann, D., Zimmermann, M. and Rösli, U., 1990. Investigation of 1987 debris flows in the Swiss Alps: general concept and geophysical soundings. Hydrology in Mountainous Regions. II-Artificial Reservoirs; Water and Slopes. Proceedings of two Lausanne Symposia, August. *IAHS Publication*, 194, 303–310.
- Haeblerli, W., Schmid, W. and Wagenbach, D., 1988. On the geometry, flow and age of firn and ice at the Colle Gnifetti core drilling site (Monte Rosa, Swiss Alps). *Zeitschrift für Gletscherkunde und Glazialgeologie*, 24 (1), 1–19.
- Haeblerli, W., Schotterer, U., Wagenbach, D., Haeblerli-Schwitter, H. and Bortenschlager, S., 1983. Accumulation characteristics on a cold, high-Alpine firn saddle from a snow-pit study on Colle Gnifetti, Monte Rosa, Swiss Alps. *Journal of Glaciology*, 29 (102), 260–271.
- Haeblerli, W., Wegmann, M. and VonderMühl, D., 1997. Slope stability problems related to glacier shrinkage and permafrost degradation in the Alps. *Eclogae Geoscientiae et Naturkatastrophen im Hochgebirge*.
- Harris, C., Arenson, L.U., Christiansen, H.H., Etzelmueller, B., Frauenfelder, R., Gruber, S., Haeblerli, W., Hauck, C., Hoelzle, M., Humlum, O., Isaksen, K., Kääb, A., Kern-Lütsch, M.A., Lehnig, M., Matsuoka, N., Murton, J.B., Nötzli, J., Phillips, M., Ross, N., Seppälä, M., Springman, S.M. and VonderMühl, D., 2009. Permafrost and climate in Europe: monitoring and modelling thermal, geomorphological and geotechnical responses. *Earth-Science Reviews*, 92, 117–171.
- Hasler, A., Gruber, S. and Haeblerli, W., 2011. Temperature variability and thermal offset in steep alpine rock and ice faces. *The Cryosphere*, 5, 977–988. doi: 10.5194/tc-5-977-2011
- Hoelzle, M. and Haeblerli, W., 1995. Simulating the effects of mean annual air temperature changes on permafrost distribution and glacier size. An example from the Upper Engadin, Swiss Alps. *Annals of Glaciology*, 21, 400–405.
- Hoelzle, M., Haeblerli, W., Dischl, M. and Peschke, W., 2003. Secular glacier mass balances derived from cumulative glacier length changes. *Global and Planetary Change*, 36 (4), 295–306.
- Hoelzle, M., Haeblerli, W. and Keller, F., 1993. Application of BTS-measurements for modelling mountain permafrost distribution. *Proceedings of the Sixth International Conference on Permafrost*, 1, 272–277.

- Huggel, C., Haeberli, W. and Käab, A., 2007. Glacial hazards – perceiving and responding to threats in four world regions. In: Orlov, E. B., Wiegandt, E. and Luckman, B.H. (eds), *Darkening Peaks – Glacial Retreat, Science and Society*. University of California Press, 68–80.
- Keller, F., 1992. Automated mapping of mountain permafrost using the program PERMAKART within the Geographical Information System ARC/INFO. *Permafrost and Periglacial Processes*, 3 (2), 133–138.
- Kieffer, H., Kargel, J.S., Barry, R., Bindshadler, R., Bishop, M., MacKinnon, D., Ohmura, A., Raup, B., Antoninetti, M., Bamber, J., Braun, M., Brown, I., Cohen, D., Copland, L., DueHagen, J., Engeset, R.V., Fitzharris, B., Fujita, K., Haeberli, W., Hagen, J.O., Hall, D., Hoelzle, M., Johansson, M., Kaeab, A., Koenig, M., Konovalov, V., Maisch, M., Paul, F., Rau, F., Reeh, N., Rignot, E., Rivera, A., de Ruyter de Wildt, M., Scambos, T., Schaper, J., Scharfen, G., Schroder, J., Solomina, O., Thompson, D. van der Veen, K., Wohlleben, T. and Young, N., 2000. New eyes in the sky measure glaciers and ice sheets. *EOS, Transactions, American Geophysical Union*, 81 (24), 265+270–271. doi:10.1029/00EO00187
- Lewkowicz, A.G., Bonnaventure, P.P., Smith, S.L. and Kuntz, Z., 2012. Spatial and thermal characteristics of mountain permafrost, northwest Canada. *Geografiska Annaler: Series A, Physical Geography*. doi:10.1111/j.1468-0459.2012.00462.x
- Linsbauer, A., Paul, F., Hoelzle, M., Frey, H. and Haeberli, W., 2009. The Swiss Alps without glaciers – a GIS-based modelling approach for reconstruction of glacier beds. *Proceedings of Geomorphometry 2009*. Zurich, Switzerland, 243–247.
- Lister, G.S., Livingstone, D.M., Ammann, B., Ariztegui, D., Haeberli, W., Lotter, A.F., Ohlendorf, C., Pfister, C., Schwander, J., Schweingruber, F., Stauffer, B. and Sturm, M., 1998. Alpine paleoclimatology. In: Cebon, P., Dahinden, U., Davies H.C., Imboden, D. and Jaeger, C.C. (eds), *Views from the Alps – Regional Perspectives on Climate Change*. MIT Press, Cambridge, 73–169.
- Naef, F., Haeberli, W. and Jäggi, M., 1989. Morphological changes in the Swiss Alps resulting from the 1987 summer storms. Hydrology of Disasters. In: Starosolszky, O. and Melder, O.M. (eds), *Proceedings of the WMO Technical Conference, Geneva, November 1988*, 36–42.
- Oerlemans, J., 2012. Linear modeling of glacier length fluctuations. *Geografiska Annaler: Series A, Physical Geography*. doi:10.1111/j.1468-0459.2012.00469.x
- Oeschger, H., Schotterer, U., Stauffer, B., Haeberli, W. and Röthlisberger, H., 1978. First results from Alpine core drilling projects. Anhang 1: ^{210}Po (^{210}Pb) dating on the Colle Gnifetti core 1976 (H. Gäggeler), Anhang 2: Sahara dust in the Alps – a short review (W. Haeberli). *Zeitschrift für Gletscherkunde und Glazialgeologie*, 13 (1/2), 193–208.
- Paul, F., Barry, R.G., Cogley, J.G., Frey, H., Haeberli, W., Ohmura, A., Ommann, C.S.L., Raup, B., Rivera, A. and Zemp, M., 2009. Recommendations for the compilation of glacier inventory data from digital sources. *Annals of Glaciology*, 50 (53), 119–126.
- Paul, F., Käab, A., Maisch, M., Kellenberger, T. and Haeberli, W., 2002. The new remote sensing-derived Swiss Glacier Inventory: I. Methods. *Annals of Glaciology*, 34, 355–361.
- Springman, S.M., Arenson, L.U., Yamamoto, Y., Maurer, H., Kos, A., Buchli, T. and Derungs, G., 2012. Multidisciplinary investigations on three rock glaciers in the Swiss Alps: legacies and future perspectives. *Geografiska Annaler: Series A, Physical Geography*. doi:10.1111/j.1468-0459.2012.00464.x
- Suter, S., Laternser, M., Haeberli, W., Frauenfelder, R. and Hoelzle, M., 2001. Cold firm and ice of high-altitude glaciers in the Alps: measurements and distribution modelling. *Journal of Glaciology*, 45 (156), 85–96.
- Townshend, J., Barry, R.G., Beran, M., Briggs, S., Caponi, C., Cilahr, J., Haeberli, W., Heal, W., Landwehr, J.M., Norse, D., Scholes, R., Solomon, A., Versegny, D., Wingham, D. and Zhao, S., 1995. *GCOS/GTOS plan for terrestrial climate-related observations; Version 1.0*. GCOS 21, WMO/TD-721, UNEP/EAP/TR/95-07.
- VonderMühl, D. and Haeberli, W., 1990. Thermal characteristics of the permafrost within an active rock glacier (Murtèl/Corvatsch, Grisons, Swiss Alps). *Journal of Glaciology*, 36 (123), 151–158.
- Wagenbach, D. and Preunkert, S., 2012. Cold, Alpine ice bodies revisited: what may we learn from their impurity and isotope content? *Geografiska Annaler: Series A, Physical Geography*. doi:10.1111/j.1468-0459.2012.00461.x
- Wegmann, M., Gudmundsson, G.H. and Haeberli, W., 1998. Permafrost changes in rock walls and the retreat of alpine glaciers: a thermal modelling approach. *Permafrost and Periglacial Processes*, 9, 23–33.
- Zimmermann, M. and Haeberli, W., 1992. Climatic change and debris flow activity in high-mountain areas – a case study in the Swiss Alps. In: Boer, M.M. and Koster, E.A. (eds), *Greenhouse-Impact on Cold-Climate Ecosystems and Landscapes*. Catena Supplement, 22, 59–72.